arXiv:astro-ph/0606398v1 16 Jun 2006

Revised: 31 May 2006

Search for compact extragalactic radio sources near massive star forming regions

Y. Xu[1,2], M. J. Reid[3], K. M. Menten[2],

and

X. W. Zheng[4]

ABSTRACT

We have used the Very Large Array to search for compact milliarcsecond-size radio sources near methanol masers in high-mass star-forming regions. Such sources are required for Very Long Baseline Interferometry phase-referencing observations. We conducted pointed observations of 234 compact sources found in the NVSS survey and find 92 sources with unresolved components and synchrotron spectral indexes. These sources are likely the cores of AGNs and, thus, good candidates for astrometric calibrators.

Subject headings: galaxies — radio continuum — astrometry

1. Introduction

Phase-referenced Very Long Baseline Interferometry (VLBI) observations can measure accurately the position of a target source relative to a reference source. This, for example, permits one to measure the trigonometric parallax and proper motions of sources in our galaxy relative to extragalactic sources. We are now carrying out a large program to do this for methanol masers in regions of high mass star formation throughout the Milky Way. Table 1 contains 38 sources showing maser emission in 12.2 GHz $2_0 - 3_{-1}E$ line of methanol that we ultimately intend to observe.

[1]Shanghai Astronomical Observatory Chinese Academy of Sciences, Shanghai 20030, China

[2]Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

[3]Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

[4]Department of Astronomy, Nanjing University Nanjing 210093, China

VLBI phase-referenced observations involve a phase calibrator and a target source. However, the distribution of known calibrators is not sufficiently dense enough to find one within $\approx 2^\circ$ of any given maser source, which is necessary for high precision astrometry: Systematic errors are usually the limiting factor in phase-referenced position measurements, and these errors scale with the angular separation between the sources. Therefore, finding calibrators as close as possible to the maser target sources is mandatory.

For this purpose, we have conducted a survey of NRAO VLA Sky Survey (NVSS; Condon et al. 1998) sources near our methanol maser targets. We chose sources whose 1.4 GHz flux densities are greater than 20 mJy and that are unresolved ($< 20''$) at the NVSS resolution. The NVSS survey typically yields about 15 such compact sources within 1° of any target position. Since our maser sources are in the Galactic plane many of these NVSS sources are compact HII regions, planetary nebulae (PNe), and, perhaps, compact supernova remnants (SNRs), and, thus, unsuitable for VLBI observations. However, some of the unresolved sources are likely extragalactic synchrotron sources.

In addition to the NVSS sources, we augmented our candidate list with sources from the literature (Gregory & Condon 1991; Ma et al. 1998) and included known compact sources as a check on our procedures. In total, we observed 234 compact sources within 1° of the maser sources.

2. Observations and Data reductions

On 2004 April 2 we made snap-shot observations of our 234 sources using the NRAO[1] VLA in its C-configuration at C-band (4.885 GHz) and U-band (14.939 GHz). Bandwidths were 50 MHz, observations were in dual circular polarization, and on-source times were between 30 and 70 seconds. The absolute flux density scale was determined from observations of 3C286 and 3C48. Occasionally we missed observations owing to electronic problems and, thus, for some sources no data could be obtained (see Table 4). The data were reduced with the NRAO Astronomical Image Processing System (AIPS) using standard procedures. We achieved rms noise levels of about 0.3 mJy at 4.885 GHz and 0.8 mJy at 14.939 GHz, respectively.

In order to obtain better positional accuracy and check on compactness, we re-observed 53 sources that had flux densities higher than 5 mJy in both, the C- and U- bands and were

[1]The National Radio Astronomy Observatory (NRAO) is operated by Associated Universities Inc., under a collaborative agreement with the U.S. National Science Foundation. Very Large Array (VLA)

unresolved in C-band. These observations were done as low-priority observations in small gaps in the VLA schedule in the A- or B-array at U-band on 2004 September 26, 27, 28, October 29, and 2005 May 1, 10, and 26, respectively. We chose U-band as it is the VLA frequency that is closest to that of the 12.2 GHz masers. The observations of each object involved 6 short (1 min duration) scans, spread over a range in time for better *uv*-coverage. To minimize the phase errors arising from temporal changes in the atmospheric phase paths, we observed a standard VLA calibrator before and after each target source. Typical map rms noise levels were about 0.2 mJy. This allowed us determine positions accurate to better than 0.1 arcsec. It is important to reach that precision for VLBI astrometry in order to minimize second-order positional errors and improve image quality.

3. Results

We detected about 92 of the 234 candidate sources with flux densities of 5 mJy at C-band with flattish or falling spectral indexes. These are likely compact extragalactic sources associated with AGN activity.

In Table 2 we list weak sources < 5 mJy at C-band observed in the C-configuration. These are too weak for current astrometric observations.

In Table 3 we list the sources that were clearly resolved. These sources are probably mostly HII regions, PNe, and SNRs. While some ultra- and hyper-compact HII regions may not be resolved by the VLA, they should have flat or rising spectra at centimeter wavelengths (Wood & Churchwell 1989; Kurtz et al. 1994), that can be easily recognized in our dual-frequency observations. Although SNRs have falling spectra, they should be resolved due to their larger size (Green 2004). The sources detected a C-band, but not at U-band, might be used as calibrators at lower frequency, e.g., for VLBI observations of 6.7 GHz CH_3OH masers, and/or 1.6 and 6.0 GHz OH masers.

Table 4 lists the parameters of the 92 sources that are unresolved at C-band, have flat or falling spectral indexes, and are potentially strong enough to be calibration sources for VLBI phase-referenced observations. Their images are shown in Figure 1 which is available in the electronic version of this paper. Except for a few sources, consisting of two components, most of the sources appear dominated by a single compact component. Most of them contain over 85% of their flux density in an unresolved component at C- and/or U-band and are likely compact extragalactic sources. Even though we selected sources with our C-configuration observations that were nearly unresolved and had > 5 mJy at C-band, when observed with the more extended A- or B-configurations some were not detected. Since these might contain

a compact component that could be used as a phase-reference calibrator at lower frequencies, we also retain them in Table 4.

Already, source 023016+620937 in Table 4 has proven to be a good calibrator for Very Long Baseline Array observations (VLBA; Xu et al. 2006). This source probably has a position uncertainty of about a few milli arcseconds (mas). Using this and two other sources we have measured a parallax (to the W3OH masers) with an accuracy of 0.007 mas. Three of our sources are incluced in the third extension to the Very Long Baseline Array (VLBI) Calibrator Survey (VCS3; Petrov et al. 2005) listed there with (sub)mas accuracies. By comparing the results from different observations listed in Table 5 we find that our C-array positions deviate from the VCS3 positions by 0.25, 0.91, and 0.14 arcsec and our A-array positions by 51 and 11 mas. We, thus, estimate that the position uncertainty is about $0.5''$ in snap-shot C-configuration observations and $0.05''$ – $0.15''$ in VLA A- and B-configuration observations, respectively, for high declination sources. Positional accuracy is likely to degrade considerably at low declinations and for weak sources (less than 10 mJy).

We calculated the two-point spectral index from 5 to 15 GHz from C-configuration observations. Most of the sources in Table 4 have a steep-spectrum with an average spectral index of $\alpha = 0.96$ ($S_\nu \sim \nu^{-\alpha}$). One must keep in mind that since these were often measured with observations within one VLA configuration, the 5 GHz beam size was about a factor of 10 larger in solid angle than the 15 GHz beam size. Thus, if a source is unresolved at 5 GHz, but partially resolved at 15 GHz, the spectral index may appear overly steep. Ulvestad et al. (1981) found a median spectral index of $\alpha = 0.09$ between 1.4 and 5 GHz for unresolved sources in a flux-limited sample. Zhang & Fan (2003) classified this sample by properties of galaxies. They found an average spectral index of 0.22 for blazers, 0.28 for QSOs, and 1.01 for galaxies, respectively. As a comparison, we obtained an average spectral index of $\alpha = 0.83$ between 1.4 and 5 GHz. Of course, since both observations spanned many years, time variability could, in principle, affect spectral index results. However general speaking, most of sources are likely to be radio galaxies.

4. Conclusions

A VLA search has resulted in the identification of 92 compact sources most likely of extragalactic nature. These sources are likely to be suitable calibrators for phase-referencing VLBI observations of 12.2 GHz CH_3OH or/and lower frequency masers.

REFERENCES

Condon, J. J., Cotton, W. D., Greisen, E. W. et al. 1998, AJ, 115, 1693

Green, D. A. 2004, A Catalogue of Galactic Supernova Remnants, Mullard Radio Astronomy Observatory, Cavendish Laboratory.

Gregory, P. C., & Condon, J. J. 1991, ApJS, 91, 1011

Kurtz, S., Churchwell, E., & Wood, D. O. S. 1994, ApJS, 91, 659

Ma, C., Arias, E. F., Eubanks, T. M., et al. 1998, AJ, 116, 516

Petrov, L., Kovalev, Y. Y., Fomalont, E., & Gordon, D., 2005, AJ, 129, 1163

Ulvestad, J., Johnston, K., Perley, R., & Fomalont, E. 1981, AJ, 86, 1010

Wood, D. O. S., & Churchwell, E. 1989, ApJS, 69, 831

Xu, Y., Reid, M. J., Zheng, X. W., & Menten, K. M. 2006, Science, 311, 54

Zhang, J. S., & Fan, J., H. 2003, ChJAA, 3, 415

This preprint was prepared with the AAS LaTeX macros v5.2.

Table 1. 12.2 GHz Methanol Maser Sources Considered

G8.68−0.37	G9.62+0.20	G10.47+0.03	G11.90−0.14	G12.03−0.03
G12.68−0.18	G12.89+0.49	G12.91−0.26	G15.03−0.68	G23.01−0.41
G23.44−0.18	G24.33+0.14	G27.36−0.16	G28.15+0.00	G29.86−0.05
G29.95−0.02	G30.20−0.17	G30.78+0.23	G30.79+0.20	G31.28+0.06
G32.74−0.08	G35.20−0.74	G35.20−1.74	G37.43+1.50	G49.49−0.37
G59.78+0.06	G109.87+2.11	G111.54+0.78	G133.95+1.07	G188.95+0.89
G232.62+0.99	G345.01+1.79[a]	G348.55−0.98	G348.70−1.04	G351.42+0.64
G353.41−0.36	G354.61+0.47	G359.61−0.24[a]		

[a]No detected compact extragalactic radio sources within 1° of this source.

Table 2. Sources with C-band Flux Densities < 5 mJy Observed in C-Configuration

172149−352620	180349−203653	181213−171318	181345−190435	181540−175738
183408−084922	184114−045503	184601−024601	184612−015715	184640−021544
184656−020132	185211+035715	185316+010040	185845+005600	192232+141303

Table 3. Sources Resolved in C- or U-Band Observations

022649+621220	022703+615232[a]	060915+223503	061032+221514	073139−161756
073223−173012	073257−164058	165451−393420	165640−401334	165646−401437[a]
171742−354155	172051−354603[a]	172052−345448	172150−364115	172201−394441
172215−382857	172804−325351	172818−350410	172827−350730	172919−343351
172928−343630	172937−351344[a]	173329−324758[a]	174451−294039	174828−293912
180416−202042	180506−193602	180513−195034[b]	180618−213731[a]	180816−171451[a]
180826−221653[b]	180827−173332	180846−192243	180849−172219	180859−200352
180939−192116[a]	181018−204544	181028−195548[a]	181031−195629[a]	181050−204110
181109−164738	181132−193042[a]	181210−184558	181345−190435[b]	181401−185324[a]
181449−173243[a]	181644−162701	181645−182524	181646−162035	181651−184129
183241−090446	183321−073121	183344−071344[a]	183432−073917	183628−070511
183657−092859	183729−082246[b]	183735−085836	183920−054509	183921−041952
183929−053553	183932−054420[a]	183955−053853[a]	184030−045722	184037−054317
184119−044418	184206−042219	184230−021839	184415−041757[a]	184600−024157
184601−010146	184621−022143	184659−020727[a]	184727−015504	184735−020143
184737−015856	184811−012630	184844−013314[a]	184916+001622[a]	184933−012907
184950−022526	185031−000155	185102−004126	185124+000412	185158+045723
185203+000926	185234+044321	185250+005527[a]	185336+042625	185343+004144[a]
185351−002508	185403+043304	185406+041156	185418+034959	185423+013529
185534+021910	185623+022328	185709+013854	185741+044212	185810+013657
185908+004150	190014+003556	190018+003719	190042−005151	190059+002757
190133+020915	190215+012223	190230−002831	190236−002657	190335+003217
192352+135603	192424+142738	192527+151310	194050+230605	194054+230544
194251+240520	194318+233032	194411+232659	194429+241937	194552+240957
225526+621739	225552+613904	225616+622227	225809+623240	231319+615644
231531+610709				

[a]Known HII region according to SIMBAD Astronomical Database and/or source with rising spectrum.

[b]Planetary nebula according to SIMBAD Astronomical Database.

Table 4. Parameters of Possible Calibrators

Source Name	Source Position		1.4 GHz	5 GHz	15 GHz	Spectral	θ	Δ
	α (2000)	δ (2000)	(mJy)	(mJy)	(mJy)	Index	($''$)	$\circ$
133.95+1.07	02 27 04.1	+61 52 22		**W3OH**				
022515+611912	02 25 15.202	+61 19 15.86	93.5	24.5			2.8	0.72
023016+620936	02 30 16.1617	+62 09 37.675	199.6	76.0	15.6	1.44	0.6[a]	0.85
188.95+0.89	06 08 53.7	+21 38 30						
060759+212945	06 07 59.5656	+21 29 43.733	45.0	17.3	6.6	1.01	0.0[b]	0.27
060831+215535	06 08 31.338	+21 55 34.59	39.2	7.3			1.6	0.30
061026+213412	06 10 26.593	+21 34 12.69	33.3	7.9			1.5	0.39
060806+211327	06 08 06.328	+21 13 26.47	32.8	6.0			1.5	0.46
060718+220452	06 07 18.656	+22 04 51.28	34.9	10.1			1.6	0.59
J0606+2159[d]	06 06 38.8201	+21 59 28.270		17.9	9.9	0.59	0.1[b]	0.66
060622+211751	06 06 22.533	+21 17 50.66	42.4	6.1			2.2	0.72
060717+221819	06 07 17.436	+22 18 18.32	67.1	31.4	7.8	0.90	1.5	0.78
J0608+2229[d]	06 08 34.327	+22 29 42.94		64.3	60.1	0.01	0.5[a]	0.85
232.62+0.99	07 32 09.6	−16 58 16						
073111−163223	07 31 11.451	−16 32 23.53	254.1	117.5	55.7	0.63	0.5[a]	0.50
073002−162904	07 30 02.496	−16 29 04.33	30.6	12.1			1.7	0.72
072924−163656	07 29 23.928	−16 36 56.07	364.1	63.5	6.2	1.96	1.0	0.78
072912−171729	07 29 12.081	−17 17 29.46	20.1	5.3			1.7	0.81
072833−164825	07 28 33.798	−16 48 26.70	39.5	12.1			1.1	0.91
072858−173000	07 28 58.175	−17 30 00.94	41.6	15.7			1.5	0.96
348.55−0.98	17 19 20.4	−39 03 51						
172216−391447	17 22 16.609	−39 14 49.39	139.0	31.6	6.2	1.56	2.1	0.76
172101−352009	17 21 01.5961	−35 20 09.121	122.2	43.2	7.7	1.55	0.4[c]	0.96
348.70-1.04	17 20 04.0	−38 58 32						
172216−391447	17 22 16.609	−39 14 49.39	139.0	31.6	6.2	1.56	2.1	0.62
172101−352009	17 21 01.5961	−35 20 09.121	122.2	43.2	7.7	1.55	0.4[c]	0.91
351.42+0.64	17 20 53.5	−35 47 02		**NGC 6334-F**				
172101−352009	17 21 01.5961	−35 20 09.121	122.2	43.2	7.7	1.55	0.4[c]	0.45
353.41−0.36	17 30 26.1	−34 41 47						
173107−343225	17 31 07.6656	−34 32 22.846	73.2	101.6	75.4	0.32	0.1[c]	0.23
173155−344959	17 31 55.4007	−34 49 59.864	265.1	89.8	22.0	1.26	0.2[c]	0.40
354.61+0.47	17 30 17.1	−33 13 55						
173157−331835	17 31 56.9970	−33 18 34.466	191.5	50.3	7.9	1.61	0.1[c]	0.42
8.68−0.37	18 06 22.9	−21 37 03						
180437−214709	18 04 38.0246	−21 47 07.564	285.6	37.5	5.2	1.92	0.2[c]	0.47
180806−212445	18 08 06.8482	−21 24 45.067	259.1	144.0	63.7	0.67	0.2[c]	0.48
J1811−2055	18 11 06.7945	−20 55 03.240			93.5		0.1[c]	1.31
9.62+0.20	18 06 14.8	−20 31 32						
180650−202742	18 06 50.2510	−20 27 40.542	113.8	59.6	5.9	1.59	0.2[b]	0.16
180323−203017	18 03 23.7239	−20 30 17.719	64.3	30.8	16.0	0.65	0.2[c]	0.71
J1811−2055	18 11 06.7945	−20 55 03.240			93.5		0.1[c]	1.21
10.47+0.03	18 08 38.2	−19 51 50						
180650−202742	18 06 50.2507	−20 27 40.537	113.8	59.6	5.9	1.59	0.2[b]	0.75
J1811−2055	18 11 06.7945	−20 55 03.240			93.5		0.1[c]	1.16
11.90−0.14	18 12 11.6	−18 41 38						

Table 4—Continued

Source Name	Source Position		1.4 GHz	5 GHz	15 GHz	Spectral	θ	Δ
	α (2000)	δ (2000)	(mJy)	(mJy)	(mJy)	Index	($''$)	$\circ$
181401−181142	18 14 01.9952	−18 11 36.524	46.5	10.7	4.4	0.81	0.1[c]	0.68
181007−181331	18 10 07.9885	−18 13 30.396	76.5	13.8	2.9	1.43	0.3[c]	0.70
181530−183614	18 15 30.3696	−18 36 13.250	71.0	64.6	56.6	0.10	0.1[c]	0.83
181445−191400	18 14 45.066	−19 13 59.50	109.3	27.3	5.2	1.12	2.0	0.84
180855−182253	18 08 55.488	−18 22 53.70	204.1	137.4	53.6	0.78	0.6[a]	0.88
180927−180451	18 09 27.878	−18 04 47.84	26.3	20.6	11.9	0.44	0.8[a]	0.92
12.03−0.03	18 12 02.5	−18 31 57						
181007−181331	18 10 07.9885	−18 13 30.396	76.5	13.8	2.9	1.43	0.3[c]	0.57
181401−181142	18 14 01.9952	−18 11 36.524	46.5	10.7	4.4	0.81	0.1[c]	0.60
180927−180451	18 09 27.878	−18 04 47.84	26.3	20.6	11.9	0.44	0.8[a]	0.79
180855−182253	18 08 55.488	−18 22 53.70	204.1	137.4	53.6	0.78	0.6[a]	0.79
181530−183614	18 15 30.3696	−18 36 13.250	71.0	64.6	56.6	0.10	0.1[c]	0.87
181445−191400	18 14 45.066	−19 13 59.50	109.3	27.3	5.2	1.12	2.0	0.98
12.68−0.18	18 13 55.1	−18 01 35						
181401−181142	18 14 01.9952	−18 11 36.524	46.5	10.7	4.4	0.81	0.1[c]	0.17
181530−183614	18 15 30.3696	−18 36 13.250	71.0	64.6	56.6	0.10	0.1[c]	0.70
181007−181331	18 10 07.9877	−18 13 30.520	76.5	13.8	0.2	1.43	0.6[c]	0.97
12.89+0.49	18 11 51.3	−17 31 29						
181134−170325	18 11 34.488	−17 03 20.54	44.9	6.2			1.7	0.47
180929−170352	18 09 28.906	−17 03 51.03	53.9	8.6			2.3	0.75
180927−180451	18 09 27.878	−18 04 47.84	26.3	20.6	11.9	0.44	0.8[a]	0.77
181007−181331	18 10 07.9885	−18 13 30.396	76.5	13.8	2.9	1.43	0.3[c]	0.82
181401−181142	18 14 01.9952	−18 11 36.524	46.5	10.7	4.4	0.81	0.1[c]	0.86
12.91−0.26	18 14 39.1	−17 52 06		**W33A**				
181401−181142	18 14 01.9952	−18 11 36.524	46.5	10.7	4.4	0.81	0.1[c]	0.36
181530−183614	18 15 30.3696	−18 36 13.250	71.0	64.6	56.6	0.10	0.1[c]	0.77
15.03−0.68	18 20 24.7	−16 11 39		**M17**				
181723−161442	18 17 22.964	−16 14 39.96	71.0	35.3	5.2	1.47	2.0	0.76
23.01−0.41	18 34 39.9	−09 00 44						
183319−085527	18 33 19.5819	−08 55 26.211	273.9	63.5	70.8	0.17	0.0[b]	0.35
183558−093724	18 35 58.7041	−09 37 29.849	144.9	18.0	3.8	1.42	0.1[c]	0.69
183424−095319	18 34 24.198	−09 53 21.49	181.2	51.1	17.2	0.79	1.3[a]	0.88
183809−084048	18 38 09.672	−08 40 41.45	66.5	7.3			2.6	0.93
23.44−0.18	18 34 39.0	−08 31 36						
183319−085527	18 33 19.5819	−08 55 26.211	273.9	63.5	70.8	0.17	0.0[b]	0.52
183809−084048	18 38 09.672	−08 40 41.45	66.5	7.3			2.6	0.89
24.33+0.14	18 35 09.4	−07 34 58						
183354−071106	18 33 54.0032	−07 11 09.436	200.0	91.5	32.2	0.95	0.1[c]	0.51
183214−070643	18 32 14.958	−07 06 45.44	99.6	18.1			3.2	0.87
183124−072551	18 31 24.746	−07 25 48.83	28.0	13.4	3.9	1.11	1.8	0.95
183758−065330	18 37 58.039	−06 53 31.00	575.7	235.5	122.6	0.48	0.6[a]	0.99
27.36−0.16	18 41 50.6	−05 01 47						
184445−045425	18 44 45.795	−04 54 23.75	107.0	24.3			1.4	0.74
28.15+0.00	18 42 40.9	−04 15 26						
184445−045425	18 44 45.795	−04 54 23.75	107.0	24.3			1.4	0.83

Table 4—Continued

Source Name	Source Position		1.4 GHz	5 GHz	15 GHz	Spectral	θ	Δ
	α (2000)	δ (2000)	(mJy)	(mJy)	(mJy)	Index	($''$)	$\circ$
184626−040853	18 46 26.874	−04 08 52.39	47.7	16.7	no data		1.7	0.95
29.86−0.05	18 45 59.9	−02 45 12						
184700−022751	18 47 00.394	−02 27 52.27	800.0	345.1	132.6	0.76	0.9[a]	0.38
184236−031108	18 42 35.941	−03 11 08.05	380.6	87.8	16.2	1.29	1.3	0.95
29.95−0.02	18 46 03.6	−02 39 24						
184700−022751	18 47 00.394	−02 27 52.27	800.0	345.1	132.6	0.76	0.9[a]	0.31
184936−021629	18 49 35.1676	−02 17 11.965		18.7	8.4	0.61	0.1[c]	0.97
30.20−0.17	18 47 02.5	−02 30 38						
184700−022751	18 47 00.394	−02 27 52.27	800.0	345.1	132.6	0.76	0.9[a]	0.05
184936−021629	18 49 35.1676	−02 17 11.965		18.7	8.4	0.61	0.1[c]	0.68
30.78+0.23	18 46 41.6	−01 48 31						
184614−011942	18 46 14.6811	−01 19 40.600	138.1	52.6	6.6	1.51	0.1[b]	0.49
184700−022751	18 47 00.394	−02 27 52.27	800.0	345.1	132.6	0.76	0.9[a]	0.66
184353−013415	18 43 53.049	−01 34 15.37	71.7	30.9			1.6	0.74
184351−012914	18 43 51.280	−01 29 17.52	371.3	52.4			1.5	0.78
184959−013256	18 49 59.3231	−01 32 57.020	1459.8	no data	47.5		0.1[b]	0.86
184936−021629	18 49 35.1676	−02 17 11.965		18.7	8.4	0.61	0.1[c]	0.86
30.79+0.20	18 46 48.2	−01 48 46						
184700−022751	18 47 00.394	−02 27 52.27	800.0	345.1	132.6	0.76	0.9[a]	0.65
184353−013415	18 43 53.049	−01 34 15.37	71.7	30.9			1.6	0.77
184351−012914	18 43 51.280	−01 29 17.52	371.3	52.4			1.5	0.81
184936−021629	18 49 35.1676	−02 17 11.965		18.7	8.4	0.61	0.1[c]	0.84
184959−013256	18 49 59.3231	−01 32 57.020	1459.8	no data	47.5		0.1[b]	0.84
31.28+0.06	18 48 12.7	−01 26 36						
184959−013256	18 49 59.3231	−01 32 57.020	1459.8	no data	47.5		0.1[b]	0.46
184614−011942	18 46 14.6811	−01 19 40.600	138.1	52.6	6.6	1.51	0.1[b]	0.51
184936−021629	18 49 35.1676	−02 17 11.965		18.7	8.4	0.61	0.1[c]	0.90
32.74−0.08	18 51 21.8	−00 12 13						
185146+003532[e]	18 51 46.7218	+00 35 32.413	839.9	771.2	625.5	0.15	0.0[b]	0.40
185343+001957	18 53 43.8523	+00 19 57.481	39.8	20.5	9.4	0.67	0.1[b]	0.61
184821+001108	18 48 21.237	+00 11 02.22	135.6	no data	8.1		0.6[a]	0.75
185152−004000	18 51 52.940	−00 40 01.85	388.8	no data	11.1		0.4[a]	0.88
35.20−0.74	18 58 13.0	+01 40 33						
190035+012119	19 00 35.444	+01 21 19.23	45.6	23.9			2.1	0.67
185749+005119	18 57 49.540	+00 51 19.29	73.6	55.3	23.8	0.82	0.5[a]	0.83
185515+021054	18 55 15.458	+02 10 50.55	84.6	19.5			1.7	0.90
185500+021544	18 55 00.116	+02 15 40.77	58.4	47.6	28.7	0.51	0.7[a]	1.00
35.20−1.74	19 01 45.5	+01 13 28						
190035+012119	19 00 35.444	+01 21 19.23	45.6	23.9			2.1	0.32
190231+004047	19 02 31.888	+00 40 47.98	31.1	10.5			1.7	0.58
190426+011036	19 04 26.3965	+01 10 38.746	61.7	63.5	48.6	-0.04	0.0[b]	0.67
190353+014526	19 03 53.0610	+01 45 26.351	196.4	137.0	72.9	0.57	0.0[b]	0.75
190227+002155	19 02 27.711	+00 21 56.69	216.1	25.9			1.4	0.88
190515+010609	19 05 15.789	+01 06 09.80	54.6	21.2			1.3	0.89
37.43+1.50	18 54 17.1	+04 41 09						

Table 4—Continued

Source Name	Source Position		1.4 GHz	5 GHz	15 GHz	Spectral	θ	Δ
	α (2000)	δ (2000)	(mJy)	(mJy)	(mJy)	Index	($''$)	$\circ$
185521+050927	18 55 21.9147	+05 09 28.468	113.1	29.8	9.5	1.08	0.1[b]	0.54
185120+050228	18 51 21.111	+05 02 35.05		7.6			1.7	0.66
185407+040059	18 54 07.9034	+04 00 59.403	126.6	40.9	14.1	0.89	0.1[c]	0.67
185618+041420	18 56 18.7659	+04 14 22.601	74.6	22.4	4.1	1.28	0.1[b]	0.68
185513+052158	18 55 13.502	+05 21 57.09	111.4	53.6	23.9	0.73	0.5[a]	0.72
185650+041520	18 56 50.269	+04 15 19.32	55.5	16.1			0.9	0.77
185124+050118	18 51 24.812	+05 01 18.84	19.9	6.5			1.2	0.79
185319+035218	18 53 19.625	+03 52 19.19	40.1	10.5			2.0	0.85
185244+053534	18 52 44.699	+05 35 35.43	53.2	6.8			2.1	0.99
49.49−0.37	19 23 39.4	+14 31 01	**W51 IRS2 (=W51d)**					
192233+150448	19 22 33.2725	+15 04 47.541	115.3	53.0	20.7	0.62	0.0[b]	0.63
59.78+0.06	19 43 10.9	+23 44 03						
194358+233010	19 43 58.1411	+23 30 10.014	118.8	34.1	15.0	0.88	0.0[b]	0.30
194313+241314	19 43 13.699	+24 13 16.50	43.4	13.3			2.0	0.49
194111+232805	19 41 11.645	+23 28 05.22	30.8	7.3			2.0	0.56
194256+230547	19 42 56.707	+23 05 49.45	20.4	8.9			1.6	0.64
194155+230755	19 41 55.1113	+23 07 57.521	70.9	32.6	8.6	1.08	0.0[b]	0.68
194619+241858	19 46 19.737	+24 18 57.60	52.5	52.9			2.2	0.98
J1946+2300[e]	19 46 06.273	+23 00 05.27		55.0	27.2	0.52	0.7[a]	1.03
109.87+2.11	22 56 18.1	+62 01 49	**Cepheus A**					
225713+621509	22 57 13.4508	+62 15 09.881	84.7	26.3	3.7	1.70	0.1[b]	0.32
J2254+6209[e]	22 54 25.2944	+62 09 38.719		58.6	24.1	0.83	0.1[b]	0.45
225452+613313	22 54 52.761	+61 33 13.43	70.6	28.8			1.6	0.59
225420+613919	22 54 20.925	+61 39 20.49	43.2	12.3			1.7	0.62
225838+611402	22 58 38.186	+61 14 03.49	42.5	9.2			1.7	0.99
111.54+0.78	23 13 45.4	+61 28 11	**NGC 7538 IRS 1**					
231310+611009	23 13 10.297	+61 10 09.37	297.3	43.5			2.1	0.33
231218+610804	23 12 18.1477	+61 08 02.072	32.0	17.9	5.3	0.92	0.1[b]	0.49
231544+615109	23 15 43.695	+61 51 18.21	32.0	11.7			2.3	0.63
231658+620347	23 16 57.747	+62 03 48.54	85.5	34.3			1.6	1.00

Note. — The first column presents source name for the masers (bold face) and the extragalactic radio sources. The next two columns list their J2000 equatorial coordinates. For the maser sources the next column gives a common name (if available). For the calibrator sources columns 4 to 6 give the integrated flux density at L- (from NVSS), peak flux density at C- and U-bands, respectively. Column 7 is the spectral index between C- and U- bands ($S_\nu \sim \nu^{-\alpha}$). Column 8 is the upper limits of source sizes (FWHM of the maximum major axes of the sources as determined from the Gaussian fits using AIPS task JMFIT). The data come from C-band observed with C-configuration, except those marked by signs. The last column gives the separation between masers and calibrators. Some sources have no data due to failure of the telescope.

[a]U-band data observed with C configuration.

[b]U-band data observed with A configuration.

[c]U-band data observed with B configuration.

[d]Source listed in Gregory & Condon (1991).

[e]Source listed in VLBA calibrator source (Petrov et al. 2005).

Table 5. Comparison of Calibrator Position Errors Derived from VLA and VLBI Data

Source Name	VLA-C Position		Δ(Y/C−VLBI)	VLA-A Position		Δ(Y/A−VLBI)	VLBI		Error
	α (2000)	δ (2000)	(arcsec)	α (2000)	δ (2000)	(arcsec)	α (2000)	δ (2000)	(mas)
023016+620936	02 30 16.068	+62 09 38.47	(−0.58, −0.75)	16.1617	37.675	(+0.076, −0.045)	16.15090	+37.7200[a]	
185146+003532	18 51 46.729	+00 35 32.59	(+0.09, +0.23)	46.7218	32.413	(−0.019, +0.050)	46.72308	+32.3628[b]	(1.43, 1.66)
J1946+2300	19 46 06.273	+23 00 05.27	(+0.30, +0.86)				06.25140	+04.4146[b]	(0.32, 0.57)
J2254+6209	22 54 25.275	+62 09 38.66	(−0.13, −0.06)	25.2944	38.719	(+0.009, −0.006)	25.29305	+38.7247[b]	(6.13, 4.93)

Note. — -Listed are, left to right, source name, position derived from VLA C-array observations, difference between the latter and the VLBI position (east offset and north offset), position derived from VLA A-array observations, difference between the latter and the VLBI position, and the VLBI position and its uncertainties. Of the coordinates for the latter two positions only seconds of time and arcseconds are given.

[a]From our VLBA observations (Xu et al. 2006).

[b]From VCS3 (Petrov et al. 2005).


6 cm C−array
022515+611912
6 cm C−array
023016+620936
2 cm C−array
023016+620936
6 cm C−array
060622+211751
2 cm A−array
J0606+2159
6 cm C−array
J0606+2159
6 cm C−array
060717+221819
2 cm C−array
060717+221819
6 cm C−array
060718+220452
2 cm A−array
060759+212945
6 cm C−array
060759+212945
6 cm C−array
060806+211327
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


6 cm C−array
060831+215535
6 cm C−array
J0608+2229
2 cm C−array
J0608+2229
6 cm C−array
061026+213412
6 cm C−array
072833−164825
6 cm C−array
072858−173000
6 cm C−array
072912−171729
6 cm C−array
072924−163656
2 cm C−array
072924−163656
6 cm C−array
073002−162904
6 cm C−array
073111−163223
2 cm C−array
073111−163223
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


2 cm B−array
172101−352009
6 cm C−array
172101−352009
6 cm C−array
172216−391447
2 cm C−array
172216−391447
2 cm B−array
173107−343225
6 cm C−array
173107−343225
2 cm B−array
173155−344959
6 cm C−array
173155−344959
2 cm B−array
173157−331835
6 cm C−array
173157−331835
2 cm B−array
180323−203017
6 cm C−array
180323−203017
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


2 cm B−array
180437−214709
6 cm C−array
180437−214709
2 cm A−array
180650−202742
6 cm C−array
180650−202742
2 cm B−arraay
180806−212445
6 cm C−array
180806−212445
6 cm C−array
180855−182253
2 cm C−array
180855−182253
6 cm C−array
180927−180451
2 cm C−array
180927−180451
6 cm C−array
180929−170352
2 cm B−array
181007−181331
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


6 cm C−array
181007−181331
2 cm B−array
J1811−2055
6 cm C−array
181134−170325
2 cm B−array
181401−181142
6 cm C−array
181401−181142
6 cm C−array
181445−191400
2 cm C−array
181445−191400
2 cm B−array
181530−183614
6 cm C−array
181530−183614
6 cm C−array
181723−161442
2 cm C−array
181723−161442
6 cm C−array
183124−072551
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


2 cm C−array
183124−072551
6 cm C−array
183214−070643
2 cm A−array
183319−085527
6 cm C−array
183319−085527
2 cm B−array
183354−−071106
6 cm C−array
183354−071106
6 cm C−array
183424−095319
2 cm C−array
183424−095319
6 cm C−array
183558−093724
6 cm C−array
183758−065330
2 cm C−array
183758−065330
6 cm C−array
183809−084048
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


6 cm C−array
184236−031108
2 cm C−array
184236−031108
6 cm C−array
184351−012914
6 cm C−array
184353−013415
6 cm C−array
184445−045425
2 cm A−array
184614−011942
6 cm C−array
184614−011942
6 cm C−array
184626−040853
6 cm C−array
184700−022751
2 cm C−array
184700−022751
2 cm C−array
184821+001108
2 cm B−array
184936−021629
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


6 cm C−array
184936−021629
2 cm B−array
184959−013256
6 cm C−array
185120+050228
6 cm C−array
185124+050118
2 cm A−array
185146+003532
6 cm C−array
185146+003532
2 cm C−array
185152−004000
6 cm C−array
185244+053534
6 cm C−array
185319+035218
2 cm A−array
185343+001957
6 cm C−array
185343+001957
2 cm B−array
185407+040059
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


6 cm C−array
185407+040059
6 cm C−array
185500+021544
2 cm C−array
185500+021544
6 cm C−array
185513+052158
2 cm C−array
185513+052158
6 cm C−array
185515+021054
2 cm A−array
185521+050927
6 cm C−array
185521+050927
2 cm A−array
185618+041420
6 cm C−array
185618+041420
6 cm C−array
185650+041520
6 cm C−array
185749+005119
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


2 cm C−array
185749+005119
6 cm C−array
190035+012119
6 cm C−array
190227+002155
6 cm C−array
190231+004047
2 cm A−array
190353+014526
6 cm C−array
190353+014526
2 cm A−array
190426+011036
6 cm C−array
190426+011036
6 cm C−array
190515+010609
2 cm A−array
192233+150448
6 cm C−array
192233+150448
6 cm C−array
194111+232805
DECLINATION (J2000)
RIGHT ASCENSION (J2000)


6 cm C−array
194155+230755
6 cm C−array
194256+230547
6 cm C−array
194313+241314
2 cm A−array
194358+233010
6 cm C−array
194358+233010
6 cm C−array
J1946+2300
2 cm C−array
J1946+2300
6 cm C−array
194619+241858
6 cm C−array
225420+613919
2 cm A−array
J2254+6209
6 cm C−array
J2254+6209
6 cm C−array
225452+613313
DECLINATION (J2000)
RIGHT ASCENSION (J2000)



Fig. 1.— Maps of the 2 and 6 cm continuum emission from extragalactic radio sources with VLA-C, -B, and -A arrays. Contour levels are -20, and 20 - 90% in steps of 10% of the peak intensity.